Fate Report for Case # P-18-0152

Fate

```
Summary Statement
                 Fate P-18-0152
           Summary FATE:
          Statement: Estimations for the low weight, monomer species, MW =
                             with \overline{MP} < 25 °C (E)
                      log Kow = -1.41 (E)
                      > 10 \text{ g/L at } 25 \text{ °C (E)}
                      VP = 1.2E-4 \text{ torr at } 25 \text{ }^{\circ}\text{C} \text{ } (E)
                      BP = 292 \, ^{\circ}C
                      (E)
                      H < 1.00E-8 (E)
                      \log Koc = 2.60 (E)
                      \log \text{ Fish BCF} = 0.50 (3)
                      (E)
                      log Fish BAF = -0.05 (1) (E)
                      POTW removal (%) = 0-25 via
                      sorption; Analog
                                                     OECD 301B(Mod Sturm CO2
                      ev):
                      3.22-6.89%/29 d. NRB.
                      Time for complete ultimate aerobic biodeg >
                      Sorption to soils/sediments = moderate
                      PBT Potential:
                      P<sub>3</sub>B<sub>1</sub>
                      *CEB FATE: Migration to ground water =
                      Bioconcentration factor to be put into E-FAST: 1
                      PMN
                      Material:
                      Overall wastewater treatment removal is 0-25% via sorption.
                      Sorption to sludge is low based on estimated physical-chemical
                      properties from EPISUITE.
                      Air Stripping (Volatilization to air) is
                      negligible based on estimated physical-chemical properties from EPISUITE.
                      Removal by biodegradation in wastewater treatment is negligible based
                                                                       : OECD 301B(Mod Sturm CO2
                      on the measured data for an analog
                      ev): 3.22-6.89%/29d NRB).
                      The aerobic aquatic biodegradation
                      half-life is greater than months based on the measured data for an analog
                                    OECD 301B(Mod Sturm CO2 ev): 3.22-6.89%/29d NRB).
                      The
                      anaerobic aquatic biodegradation half-life is greater than months based on
```

the aerobic biodegradation half-life. The anaerobic biodegradation half-life is projected to be greater than or equal to the aerobic biodegradation half-life.

Sorption to soil and sediment is moderate

based on estimated physical-chemical properties from EPISUITE.

Migration to groundwater is moderate based on estimated physical-chemical properties from EPISUITE.

PMN Material:

High

Persistence (P3) is based on the aerobic and anaerobic biodegradation half-lives.

Low Bioaccumulation potential (B1) is based on BCFBAF model estimates.

Bioconcentration/Bioaccumulation factor to be put into E-Fast: 1.

CBI:
Fate Lee, WenHsiung
Assessor:
SMILES:

Physical Properties

Property	Measured/Calculated Value	EPI
Molecular Form:		
Molecular Wt.:		
% < 500:		_
% < 1000:		

Property	Measured	Method	Estimated	Method	EPI
	Value		Value		
Melting					NaN
Point:					°C (Exp.)
					282.0794982910156 °C (Est.,
					Joback)
					119.01715087890625
					°C (Est., Gold)
					151.62960815429688 °C (Est.,
					Selected)
					1.9112759042637781%
Boiling Point:			327		NaN
					°C (Exp.)
					398.49 °C (Est.)

Proporty	Measured	Method	Estimated	Method	EPI
Property	Value	Method	Value	Method	EFI
	, and		7 mac		671.6512451171875 °K
					(Est.)
BP Pressure:					,
Vapor			0.000012		NaN
Pressure:					mmHg (Exp.)
					8.776204236904395E-8 Pa
					(Est., Antoine)
					6.582712708258498E-10
					mmHg (Est., Antoine)
					5.925547576351211E-7 Pa
					(Est., Grain)
					4.444538468033191E-9 mmHg
					(Est., Grain)
					1.3811481717325024E-4 Pa
					(Est., Mackay)
					1.0359491844800576E-6
					mmHg
					(Est., Mackay)
					5.925547576351211E-7 Pa
					(Est., Selected)
					4.444538468033191E-9 mmHg
					(Est., Selected)
					1.155788608437547E-5 Pa
					(Est., SubCooled)
					8.669151441154099E-8 mmH
					(Est.,
Water			1000		SubCooled) NaN
Solubility:			1000		(Exp.)
Solubility.					3518.090087890625 (Est.)
Log			-1.41		22 20.03 000 7 00 20 (20.0)
P:					
Log					NaN
Kow:					(Exp.)
					1.21 (Est.)
Log	NaN				12.086405502072507 (Est., log
Koc:					(MCI))
					2.7278528408647205 (Est., log
					(Kow))
					177443.140625 L/kg (Est.,
					MCI)
					15.300000190734863 L/kg
					(Est., Kow)
					(250, 11011)

Method	Estimated	Method	EPI
	Value		
			3.16 L/kg wet-wt
			NaN atm-m3/mole (Exp.)
			2.6262408007069256E-19 atm-
			m3/mole (Est., Bond)
			0.0 atm-m3/mole
			(Est., Group)
		Value	Value

pH: pH Comment:

Fate Analysis

Hydrolysis	Ve	olatilization		Volatilization
(t1/2,		(t1/2)		(t1/2)
da):	_	River (hr):		- Lake (da):
Atm Ox	0.05200538106753519	Atm Ox	0.0	Atm Ox
Potential		Potential		Potential
(t1/2)OH		(t1/2)O3		(t1/2) Total
(hr):		(hr):		(hr):
MITI	0.30000001192092896	MITI	0.0	
Linear:]	NonLinear:		
Biodeg	1.100000023841858	Biodeg	0.8999999761581421	
Linear:]	NonLinear:		
Biodeg	2.5999999046325684	Biodeg	3.5	
Survey	Sı	urvey Prim:		
ult:				
STP (%	1.9112759042637781%	STP (%	0.092710892425102%	
removal)		removal)		
Total:		Biodeg:		
STP (%	1.818569329539617%	STP (%	1.4935728189854662E-	
removal)		removal) i	14%	
Ads:		Air:		

Rationales

Removal in
Wastewater
Treatment:
Atmospheric
Oxidation:
Hydrolysis:
Photolysis:
Aerobic
Biodegradation:

Anaerobic **Biodegradation:** Sorption to Soil and **Sediment:** Migration to Groundwater: Persistence - Air: Persistence - Water: Volatilization from Water: Soil: **Sediment:** Other: Standard: Bioaccumulation:

PBT Ratings

Persistence	Bioaccumulation	Toxicity	PBT Comments
3	1		

Exposure-Based Testing

Exposure-Based
Testing:

Fate Ratings Removal in WWT/POTW

(Overall):

Removal in 0-25 WWT/POTW (Overall):

Condition	Rating Values	Rating Description				Comment
		1	2	3	4	
WWT/POTW Sorption:	1	Low	Moderate	Strong	V. Strong	
WWT/POTW Stripping:	4	Extensive	Moderate	Low	Negligible	
Biodegradation Removal:	4	Unknown	High	Moderate	Negligible	
Biodegradation Destruction:		Unknown	Complete	Partial	_	
	4		Weeks	Months		

Condition	Rating Values	Rating Description				Comment
		1	2	3	4	
Aerobic		<=			>	
Biodeg Ult:		Days			Months	
Aerobic Biodeg		<= Days	Weeks	Months	>	
Prim:					Months	
Anaerobic Biodeg	4	<= Days	Weeks	Months	>	
Ult:					Months	
Anaerobic Biodeg		<= Days	Weeks	Months	>	
Prim:					Months	
Hydrolysis (t1/2 at		<=	Hours	Days	>=	
pH		Minutes			Months	
7,25C) A:			**			
Hydrolysis (t1/2 at		<=	Hours	Days	>=	
pH		Minutes			Months	
7,25C) B:		V.	Cteana	Moderate	Low	
Sorption to Soils/Sediments:	3		Strong	Moderate	Low	
Migration to	3	Strong	Slow	Moderate	Danid	
Ground Water:	3	Negligible	Slow	Moderate	Rapid	
Photolysis A, Direct:		Negligible	Slow	Moderate	Rapid	
Photolysis B,		Negligible	Slow	Moderate	Rapid	
Indirect:						
Atmospheric Ox A, OH:		Negligible	Slow	Moderate	Rapid	
Atmospheric Ox B, O3:		Negligible	Slow	Moderate	Rapid	

Bio

Comments:

Bio The EPI outpurt file for
Comments: the PMN material with manually entered properties is attached.

Composition by

The submitter also stated that the product distribution depended on the

A fate study summary is available. Fish log BAF = -0.05 (1).

Fate Comments:

Fate Comments: Analog ev): OECD 301B(Mod Sturm CO2 ev): 3.22-6.89%/29 d. NRB.

Comments/Telephone

Artifact	Update/Upload Time	